

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 May 2001 (17.05.2001)

PCT

(10) International Publication Number  
**WO 01/34585 A1**

(51) International Patent Classification<sup>7</sup>: C07D 277/36,  
417/12, 403/12, A61K 31/426, 31/427, 31/4178, A61P  
7/00

[ES/US]; 701 Pondview Drive, Audubon, PA 19403 (US).  
DUFFY, Kevin, J. [GB/US]; 712 Mill Grove Drive, Nor-  
ristown, PA 19403 (US). SHAW, Antony [GB/US]; 713  
Jacques Circle, Chester Springs, PA 19425 (US).

(21) International Application Number: PCT/US00/30383

(74) Agents: DUSTMAN, Wayne, J. et al.; SmithKline  
Beecham Corporation, Corporate Intellectual Property,  
UW2220, 709 Swedeland Road, PO Box 1539, King of  
Prussia, PA 19406-0939 (US).

(22) International Filing Date:  
3 November 2000 (03.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/163,907 5 November 1999 (05.11.1999) US

(81) Designated States (*national*): AU, CA, JP, US.

(84) Designated States (*regional*): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).


(71) Applicant (*for all designated States except US*):  
SMITHKLINE BEECHAM CORPORATION  
[US/US]; One Franklin Plaza, Philadelphia, PA 19103  
(US).

Published:  
— With international search report.


(72) Inventors; and  
(75) Inventors/Applicants (*for US only*): LUENGO, Juan, I.

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: SEMICARBAZONE DERIVATIVES AND THEIR USE AS THROMBOPOIETIN MIMETICS



(I)



(Ia)

(57) Abstract: Compounds of  
formula (I), particularly a compound  
of formula (Ia), are non-peptide TPO  
mimetics, useful in the treatment of  
thrombocytopenia.

WO 01/34585 A1

## SEMICARBAZONE DERIVATIVES AND THEIR USE AS TROMBOPOIETIN MIMETICS

### FIELD OF THE INVENTION

5 This invention relates to thrombopoietin (TPO ) mimetics and their use as promoters of thrombopoiesis and megakaryocytopoiesis.

### BACKGROUND OF THE INVENTION

10 Megakaryocytes are bone marrow-derived cells, which are responsible for producing circulating blood platelets. Although comprising <0.25% of the bone marrow cells in most species, they have >10 times the volume of typical marrow cells. See Kuter et al. Proc. Natl. Acad. Sci. USA 91: 11104-11108 (1994). Megakaryocytes undergo a process known as endomitosis whereby they replicate their nuclei but fail to undergo cell division and thereby give rise to polyploid cells. In response to a decreased platelet count, 15 the endomitotic rate increases, higher ploidy megakaryocytes are formed, and the number of megakaryocytes may increase up to 3-fold. See Harker J. Clin. Invest. 47: 458-465 (1968). In contrast, in response to an elevated platelet count, the endomitotic rate decreases, lower ploidy megakaryocytes are formed, and the number of megakaryocytes may decrease by 50%.

20 The exact physiological feedback mechanism by which the mass of circulating platelets regulates the endomitotic rate and number of bone marrow megakaryocytes is not known. The circulating thrombopoietic factor involved in mediating this feedback loop is now thought to be thrombopoietin (TPO). More specifically, TPO has been shown to be the main humoral regulator in situations involving thrombocytopenia. See, e.g., Metcalf Nature 369:519-520 25 (1994). TPO has been shown in several studies to increase platelet counts, increase platelet size, and increase isotope incorporation into platelets of recipient animals. Specifically, TPO is thought to affect megakaryocytopoiesis in several ways: (1) it produces increases in megakaryocyte size and number; (2) it produces an increase in DNA content, in the form of polyploidy, in megakaryocytes; (3) it increases megakaryocyte endomitosis; (4) it produces 30 increased maturation of megakaryocytes; and (5) it produces an increase in the percentage of precursor cells, in the form of small acetylcholinesterase-positive cells, in the bone marrow.

Because platelets (thrombocytes) are necessary for blood clotting and when their numbers are very low a patient is at risk of death from catastrophic hemorrhage, TPO has potential useful application in both the diagnosis and the treatment of various 35 hematological disorders, for example, diseases primarily due to platelet defects. Ongoing clinical trials with TPO have indicated that TPO can be administered safely to patients. In addition, recent studies have provided a basis for the projection of efficacy of TPO therapy

in the treatment of thrombocytopenia, and particularly thrombocytopenia resulting from chemotherapy, radiation therapy, or bone marrow transplantation as treatment for cancer or lymphoma. See e.g., McDonald (1992) Am. J. Ped. Hematology/Oncology 14: 8-21 (1992).

5       The gene encoding TPO has been cloned and characterized. See Kuter et al., Proc. Natl. Acad. Sci. USA 91: 11104-11108 (1994); Barley et al., Cell 77: 1117-1124 (1994); Kaushansky et al., Nature 369:568-571 (1994); Wendling et al., Nature 369: 571-574 (1994); and Sauvage et al., Nature 369: 533-538 (1994).  
10       Thrombopoietin is a glycoprotein with two distinct regions separated by a potential Arg-Arg cleavage site. The amino-terminal region is highly conserved in man and mouse, and has some homology with erythropoietin and interferon-alpha and interferon-beta. The carboxy-terminal region shows wide species divergence.

      The DNA sequences and encoded peptide sequences for human TPO receptor (TPO-R; also known as c-mpl) have been described. See, Vigon et al. Proc. Natl. Acad. Sci. USA  
15       89: 5640-5644 (1992). TPO-R is a member of the haematopoietin growth factor receptor family, a family characterized by a common structural design of the extracellular domain, including for conserved C residues in the N-terminal portion and a WSXWS motif close to the transmembrane region. See Bazan Proc. Natl. Acad. Sci. USA 87: 6934-6938 (1990).  
20       Evidence that this receptor plays a functional role in hematopoiesis includes observations that its expression is restricted to spleen, bone marrow, or fetal liver in mice (see Souyri et al. Cell 63: 1137-1147 (1990)) and to megakaryocytes, platelets, and CD34<sup>+</sup> cells in humans (see Methia et al. Blood 82: 1395-1401 (1993)). Further evidence for TPO-R as a  
25       key regulator of megakaryopoiesis is the fact that exposure of CD34<sup>+</sup> cells to synthetic oligonucleotides antisense to TPO-R RNA significantly inhibits the appearance of megakaryocyte colonies without affecting erythroid or myeloid colony formation. Some  
30       workers postulate that the receptor functions as a homodimer, similar to the situation with the receptors for G-CSF and erythropoietin.


      The slow recovery of platelet levels in patients suffering from thrombocytopenia is a serious problem, and has lent urgency to the search for a blood growth factor agonist able to  
30       accelerate platelet regeneration.

      It would be desirable to provide compounds which allow for the treatment of thrombocytopenia by acting as a TPO mimetic.

      As disclosed herein it has unexpectedly been discovered that certain substituted thiosemicarbazone derivatives are effective as agonists of the TPO receptor, they are potent  
35       TPO mimetics.

SUMMARY OF THE INVENTION

This invention relates to compounds of Formula (I):




5 wherein:

R<sup>1</sup> and R<sup>2</sup> are each independently selected from hydrogen, C<sub>1-12</sub>alkyl, aryl, substituted aryl, and alkyl substituted with one or more substituents selected from the group consisting of: alkoxy, acyloxy, aryl, substituted aryl, amino, N-acylamino, oxo, hydroxy, cycloalkyl, substituted cycloalkyl, -C(O)OR<sup>7</sup>, -S(O)<sub>2</sub>NR<sup>7</sup>R<sup>8</sup>, -S(O)<sub>n</sub>R<sup>6</sup>, aryloxy, nitro, cyano, halogen, and protected -OH, where

R<sup>6</sup> is selected from hydrogen, alkyl, cycloalkyl, C<sub>1-C12</sub>aryl, substituted alkyl, substituted cycloalkyl and substituted C<sub>1-C12</sub>aryl, and


R<sup>7</sup> and R<sup>8</sup> are independently selected from hydrogen, cycloalkyl, aryl, substituted cycloalkyl, substituted aryl, alkyl or alkyl substituted with one or more substituents selected from the group consisting of: alkoxy, acyloxy, aryloxy, amino, N-acylamino, oxo, hydroxy, -C(O)OR<sup>6</sup>, -S(O)<sub>n</sub>R<sup>6</sup>, -C(O)NR<sup>6</sup>R<sup>6</sup>, -S(O)<sub>2</sub>NR<sup>6</sup>R<sup>6</sup>, nitro, cyano, cycloalkyl, substituted cycloalkyl, halogen, C<sub>1-C12</sub>aryl, substituted C<sub>1-C12</sub>aryl and protected -OH where R<sup>6</sup> is as described above; and

n is 0-3; or R<sup>1</sup> and R<sup>2</sup> taken together with the



group to which they are

attached represent a ring of formula (A):




where  $R^4$  and  $R^{10}$  are each independently selected from two hydrogens,  $=NR^5$ ,  $=O$ ,  $=S$ , and  $=CHR^5$ , where  $R^5$  is  $C_1$ - $C_{12}$ aryl or substituted  $C_1$ - $C_{12}$ aryl; and  $m$  is 0 to 2;

5 Z is a bond or selected from S or  $NR^5$ , where  $R^5$  is  $C_1$ - $C_{12}$ aryl or substituted  $C_1$ - $C_{12}$ aryl;

$R^3$  is selected from hydrogen,  $C_1$ - $C_{10}$ alkyl, phenyl, substituted phenyl, carboxyl or  $C_1$ - $C_{10}$ alkoxycarbonyl;

10

L is a group of formula (L):



(L)

15 where A, B, D and E independently represent  $CR^{11}$  or N; where  $R^{11}$  is selected from hydrogen, halogen,  $-CF_3$ ,  $-CN$ ,  $-SO_3H$ ,  $-SO_3Na$ ,  $-SO_2R^{14}$ ,  $-NO_2$ , phenyl, substituted phenyl,  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_{10}$ alkoxy,  $C_1$ - $C_{10}$ acyloxy, arylalkoxy,  $-COR^{14}$ ,  $-NR^{12}R^{13}$ , hydroxy or cycloalkyl;

where  $R^{14}$  is selected from hydroxy,  $C_1$ - $C_{10}$ alkyl, phenyl, amino, mono- or dialkylamino;

20

$R^{12}$  and  $R^{13}$  are independently selected from hydrogen,  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_{12}$ aryl, substituted  $C_1$ - $C_{12}$ aryl,  $C_1$ - $C_{10}$ acyl or cycloalkyl;

or either  $A=B$  or  $D=E$  alternatively represent O, S or  $NR^{12}$ ; where  $R^{12}$  is as defined above;

25

Y is selected from  $-S$ ,  $-O$  and  $-NR^{15}$ , where  $R^{15}$  is selected from hydrogen,  $C_1$ - $C_{10}$ alkyl, substituted  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_6$ alkylphenyl, substituted  $C_1$ - $C_6$ alkylphenyl,  $C_1$ - $C_{10}$ acyl, substituted  $C_1$ - $C_{10}$ acyl, or  $SO_2R^9$ , where  $R^9$  is  $C_1$ - $C_{10}$ alkyl, substituted  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_{12}$ aryl or substituted  $C_1$ - $C_{12}$ aryl; and

30

X is selected from  $-SR^{16}$ ,  $-OR^{16}$  or  $-NHR^{17}$ ; where  $R^{16}$  is hydrogen,  $C_1$ - $C_{10}$ alkyl or substituted  $C_1$ - $C_{10}$ alkyl;

$R^{17}$  is hydrogen,  $C_1$ - $C_{10}$ alkyl, substituted  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_6$ alkylphenyl,  $C_1$ - $C_{10}$ acyl, substituted  $C_1$ - $C_{10}$ acyl or  $SO_2R^9$ ; where  $R^9$  is  $C_1$ - $C_{10}$ alkyl, substituted  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_{12}$ aryl or substituted  $C_1$ - $C_{12}$ aryl; and

pharmaceutically acceptable salts, hydrates, solvates and esters thereof,

provided that;

5

when  $R^1$  and  $R^2$  do not form a ring and X is not  $-NHSO_2R^9$ ,  $R^5$  is not a substituted or unsubstituted pyridyl or a substituted or unsubstituted phenyl.

10 This invention relates to a method of treating thrombocytopenia, which comprises administering to a subject in need thereof an effective amount of a TPO mimetic compound of Formula (I).

The present invention also relates to the discovery that the compounds of Formula (I) are active as agonists of the TPO receptor.

15

In a further aspect of the invention there is provided novel processes and novel intermediates useful in preparing the presently invented TPO mimetic compounds.

20 Included in the present invention are pharmaceutical compositions comprising a pharmaceutical carrier and compounds useful in the methods of the invention.

Also included in the present invention are methods of co-administering the presently invented TPO mimetic compounds with further active ingredients.

25


### DETAILED DESCRIPTION OF THE INVENTION


This invention relates to compounds of Formula (I) as described above.

30 Preferred among the presently invented Formula I compounds are those in which  $R^5$  is  $C_1$ - $C_{12}$ aryl substituted with a carboxy or sulfonic acid substituent.

Preferred among the presently invented Formula I compounds are those in which  $R^1$  and  $R^2$  are selected from hydrogen,  $C_1$ - $10$ alkyl, benzyl, substituted benzyl, phenyl,

35 substituted phenyl, or  $R^1$  and  $R^2$  taken together with the attached represent a ring of formula (A):





(A)

where R<sup>4</sup> and R<sup>10</sup> are each independently selected from two hydrogens, =O, or =CHR<sup>5</sup>, where R<sup>5</sup> is C<sub>1</sub>-C<sub>12</sub>aryl or substituted C<sub>1</sub>-C<sub>12</sub>aryl; and m is 0 to 2.

5

Preferred among the presently invented Formula I compounds are those in which:

Z is S or -NR<sup>5</sup> where R<sup>5</sup> is phenyl substituted with a carboxy or sulfonic acid substituent, a six membered aromatic ring containing from 1 to 3 heteroatoms and substituted with a carboxy or sulfonic acid substituent, or a C<sub>1</sub>-C<sub>2</sub>alkylphenyl substituted with a carboxy or sulfonic acid substituent;

10

L is C<sub>3</sub>-C<sub>6</sub>aryl optionally substituted with from 1 to 3 substituents selected from the group consisting of: Br, Cl, CF<sub>3</sub>, F, -CH<sub>3</sub> and substituted phenyl;

15

Y is S; and

X is -OH; and

20

pharmaceutically acceptable salts, hydrates, solvates and esters thereof,

provided that;

25

when R<sup>1</sup> and R<sup>2</sup> do not form a ring, R<sup>5</sup> is not a substituted or unsubstituted pyridyl or a substituted or unsubstituted phenyl.

Preferred among the presently invented compounds are:

3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one  
(Compound A);

30 3-(3-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1H-pyrazol-4-yl)methylene]amino]-2-thioxoimidazolidin-4-one;

- 3-(4-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one;  
5-(4-carboxybenzylidene)-3-[(1-(3,4-dimethylphenyl)-4-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one; and  
5 5-(3-carboxybenzylidene)-3-[(1-(3,4-dimethylphenyl)-4-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one.

Compounds of Formula (I) are included in the pharmaceutical compositions of the invention and used in the methods of the invention.

10

By the term "protected hydroxy" or "protected -OH" as used herein, is meant the alcoholic or carboxylic-OH groups which can be protected by conventional blocking groups in the art as described in "Protective Groups In Organic Synthesis" by Theodora W. Greene, Wiley-Interscience, 1981, New York. Compounds containing protected hydroxy groups  
15 may also be useful as intermediates in the preparation of the pharmaceutically active compounds of the invention.

By the term "aryl" as used herein, unless otherwise defined, is meant a cyclic or polycyclic aromatic ring containing from 1 to 14 carbon atoms and optionally containing from one to five heteroatoms, provided that when the number of carbon atoms is 1 the  
20 aromatic ring contains at least four heteroatoms, when the number of carbon atoms is 2 the aromatic ring contains at least three heteroatoms, when the number of carbons is 3 the aromatic ring contains at least two heteroatoms and when the number of carbon atoms is 4 the aromatic ring contains at least one heteroatom.

By the term "C<sub>1</sub>-C<sub>12</sub>aryl" as used herein, unless otherwise defined, is meant  
25 phenyl, naphthalene, 3,4-methylenedioxyphenyl, pyridine, biphenyl, quinoline, pyrimidine, quinazoline, thiophene, furan, pyrrole, pyrazole, imidazole and tetrazole.

By the term "C<sub>3</sub>-C<sub>6</sub>aryl" as used herein, unless otherwise defined, is meant a cyclic or polycyclic aromatic ring containing from 3 to 6 carbon atoms and optionally containing from one to 4 heteroatoms, provided that when the number of carbon atoms is 3 the  
30 aromatic ring contains at least two heteroatoms and when the number of carbon atoms is 4 the aromatic ring contains at least one heteroatom.

By the term "substituted" as used herein, unless otherwise defined, is meant that the subject chemical moiety has one or more substituents selected from the group consisting of: hydroxyalkyl, alkoxy, acyloxy, alkyl, aryl, amino, N-acylamino, hydroxy,  
35 -(CH<sub>2</sub>)<sub>g</sub>C(O)OR<sup>6</sup>, -S(O)<sub>n</sub>R<sup>7</sup>, nitro, cyano, halogen, trifluoromethyl and protected -OH, where g is 0-6, R<sup>6</sup> is hydrogen or alkyl, n is 0-3, and R<sup>7</sup> is hydrogen or alkyl.



By the term "alkoxy" as used herein is meant -Oalkyl where alkyl is as described herein including -OCH<sub>3</sub> and -OC(CH<sub>3</sub>)<sub>2</sub>CH<sub>3</sub>.

The term "cycloalkyl" as used herein unless otherwise defined, is meant a nonaromatic, unsaturated or saturated, cyclic or polycyclic C<sub>3</sub>-C<sub>12</sub>.

5 Examples of cycloalkyl and substituted cycloalkyl substituents as used herein include: cyclohexyl, 4-hydroxy-cyclohexyl, 2-ethylcyclohexyl, propyl 4-methoxycyclohexyl, 4-methoxycyclohexyl, 4-carboxycyclohexyl and cyclopentyl.

By the term "acyloxy" as used herein is meant -OC(O)alkyl where alkyl is as described herein. Examples of acyloxy substituents as used herein include: -OC(O)CH<sub>3</sub>, -  
10 OC(O)CH(CH<sub>3</sub>)<sub>2</sub> and -OC(O)(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub>.

By the term "N-acylamino" as used herein is meant -N(H)C(O)alkyl, where alkyl is as described herein. Examples of N-acylamino substituents as used herein include: -N(H)C(O)CH<sub>3</sub>, -N(H)C(O)CH(CH<sub>3</sub>)<sub>2</sub> and -N(H)C(O)(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub>.

By the term "aryloxy" as used herein is meant -OC<sub>6</sub>-C<sub>12</sub>aryl where C<sub>6</sub>-C<sub>12</sub>aryl is  
15 phenyl, naphthyl, 3,4-methylenedioxyphenyl, pyridyl or biphenyl optionally substituted with one or more substituents selected from the group consisting of: alkyl, hydroxyalkyl, alkoxy, trifluoromethyl, acyloxy, amino, N-acylamino, hydroxy, -(CH<sub>2</sub>)<sub>g</sub>C(O)OR<sup>6</sup>, -S(O)<sub>n</sub>R<sup>7</sup>, nitro, cyano, halogen and protected -OH, where g is 0-6, R<sup>6</sup> is hydrogen or alkyl, n is 0-3 and R<sup>7</sup> is hydrogen or alkyl. Examples of aryloxy substituents as used herein  
20 include: phenoxy, 4-fluorophenyloxy and biphenyloxy.

By the term "heteroatom" as used herein is meant oxygen, nitrogen or sulfur.

By the term "halogen" as used herein is meant a substituent selected from bromide, iodide, chloride and fluoride.

By the term "alkyl" and derivatives thereof and in all carbon chains as used herein  
25 is meant a linear or branched, saturated or unsaturated hydrocarbon chain having C<sub>1</sub>-C<sub>12</sub> carbon atoms. Examples of alkyl substituents as used herein include: -CH<sub>3</sub>, -CH<sub>2</sub>-CH<sub>3</sub>, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -C(CH<sub>3</sub>)<sub>3</sub>, -(CH<sub>2</sub>)<sub>3</sub>-CH<sub>3</sub>, -CH<sub>2</sub>-CH(CH<sub>3</sub>)<sub>2</sub> and -CH(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>3</sub>, -CH=CH<sub>2</sub>.

By the term "treating" and derivatives thereof as used herein, is meant prophylactic  
30 or therapeutic therapy.


All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as though fully set forth.

Compounds of Formula (I) are included in the pharmaceutical compositions of the invention and used in the methods of the invention. Where a -COOH or -OH group is  
35 present, pharmaceutically acceptable esters can be employed, for example methyl, ethyl, pivaloyloxymethyl, and the like for -COOH, and acetate maleate and the like for -OH, and

those esters known in the art for modifying solubility or hydrolysis characteristics for use as sustained release or prodrug formulations.

The novel compounds of Formula I are prepared as shown in Scheme I below wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, Z, Y, L and X are as defined in Formula I and provided that these substituents do not include any such substituents that render inoperative the Scheme I process. All of the starting materials are commercially available or are readily made from commercially available starting materials by those of skill in the art.

Scheme 1



Compounds 1, are condensed with carbonyl compounds 2, available commercially or prepared by literature methods, in a suitable solvent with or without the addition of an acid catalyst such as HCl to furnish the final compound 3.

The treatment of thrombocytopenia, as described herein, is accomplished by enhancing the production of platelets.

By the term "co-administering" and derivatives thereof as used herein is meant either simultaneous administration or any manner of separate sequential administration of a TPO mimetic compound, as described herein, and a further active ingredient or ingredients, known to treat thrombocytopenia, including chemotherapy-induced thrombocytopenia and bone marrow transplantation and other conditions with depressed platelet production. Preferably, if the administration is not simultaneous, the compounds are administered in a close time proximity to each other. Furthermore, it does not matter if the compounds are administered in the same dosage form, e.g. one compound may be administered topically and another compound may be administered orally.

Because the pharmaceutically active compounds of the present invention are active as TPO mimetics they exhibit therapeutic utility in treating thrombocytopenia and other conditions with depressed platelet production.

In determining potency as TPO mimetics, the following assays were employed:

Luciferase Assay

Compounds of the present invention were tested for potency as mimetics of the TPO receptor in a Luciferase assay such as described in Lamb, et al., Nucleic Acids Research 23: 3283-3289 (1995) and Seidel, et al., Proc. Natl. Acad. Sci., USA 92: 3041-3045 (1995) by substituting a TPO-responsive BaF3 cell line (Vigon et al. Proc. Natl. Acad. Sci. USA 1992, 89, 5640-5644) for the HepG2 cells utilized therein. The murine BaF3 cells express TPO receptors and closely match the pattern of STAT (signal transducers and activators of transcription) activation observed in primary murine and human bone marrow cells in response to TPO.

Some of the preferred compounds of this invention were also active in an in vitro proliferation assay using the murine 32D-mpl cell line (Bartley, T. D. et al., Cell, 1994, 77, 1117-1124). 32D-mpl cells express Tpo-R and their survival is dependent on the presence of TPO.

The pharmaceutically active compounds within the scope of this invention are useful as TPO mimetics in mammals, including humans, in need thereof.

Within the scope of the invention Compound A showed activation of about 9% of control (control is the maximal response to TPO) at a concentration of 10 uM in the luciferase assay.

Some of the preferred compounds within the scope of the invention showed activation from about 0% to 9% control at a concentration of 1-10 uM in the luciferase assay. The preferred compounds of the invention also promoted the proliferation of 32D-mpl cells at a concentration of 10 to 30 uM.

The present invention therefor provides a method of treating thrombocytopenia and other conditions with depressed platelet production, which comprises administering a compound of Formula (I), and pharmaceutically acceptable salts, hydrates, solvates and esters thereof in a quantity effective to enhance platelet production. The compounds of Formula (I) also provide for a method of treating the above indicated disease states because of their demonstrated ability to act as TPO mimetics. The drug may be administered to a patient in need thereof by any conventional route of administration, including, but not limited to, intravenous, intramuscular, oral, subcutaneous, intradermal, and parenteral.

The pharmaceutically active compounds of the present invention are incorporated into convenient dosage forms such as capsules, tablets, or injectable preparations. Solid or liquid pharmaceutical carriers are employed. Solid carriers include, starch, lactose, calcium sulfate dihydrate, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, and stearic acid. Liquid carriers include syrup, peanut oil, olive oil, saline, and water. Similarly, the carrier or diluent may include any prolonged release material, such as glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of solid

carrier varies widely but, preferably, will be from about 25 mg to about 1 g per dosage unit. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampoule, or an aqueous or nonaqueous liquid suspension.

5           The pharmaceutical preparations are made following conventional techniques of a pharmaceutical chemist involving mixing, granulating, and compressing, when necessary, for tablet forms, or mixing, filling and dissolving the ingredients, as appropriate, to give the desired oral or parenteral products.

10           Doses of the presently invented pharmaceutically active compounds in a pharmaceutical dosage unit as described above will be an efficacious, nontoxic quantity preferably selected from the range of 0.001 - 100 mg/kg of active compound, preferably 0.001 - 50 mg/kg. When treating a human patient in need of a TPO mimetic, the selected dose is administered preferably from 1-6 times daily, orally or parenterally. Preferred forms of parenteral administration include topically, rectally, transdermally, by injection  
15 and continuously by infusion. Oral dosage units for human administration preferably contain from 0.05 to 3500 mg of active compound. Oral administration, which uses lower dosages is preferred. Parenteral administration, at high dosages, however, also can be used when safe and convenient for the patient.

20           Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular TPO mimetic in use, the strength of the preparation, the mode of administration, and the advancement of the disease condition. Additional factors depending on the particular patient being treated will result in a need to adjust dosages, including patient age, weight, diet, and time of administration.

25           The method of this invention of inducing TPO mimetic activity in mammals, including humans, comprises administering to a subject in need of such activity an effective TPO mimetic amount of a pharmaceutically active compound of the present invention.

          The invention also provides for the use of a compound of Formula (I) in the manufacture of a medicament for use as a TPO mimetic.

30           The invention also provides for the use of a compound of Formula (I) in the manufacture of a medicament for use in therapy.

          The invention also provides for the use of a compound of Formula (I) in the manufacture of a medicament for use in enhancing platelet production.

          The invention also provides for the use of a compound of Formula (I) in the manufacture of a medicament for use in treating thrombocytopenia.

35           The invention also provides for a pharmaceutical composition for use as a TPO mimetic which comprises a compound of Formula (I) and a pharmaceutically acceptable carrier.

The invention also provides for a pharmaceutical composition for use in the treatment of thrombocytopenia which comprises a compound of Formula (I) and a pharmaceutically acceptable carrier.

5 The invention also provides for a pharmaceutical composition for use in enhancing platelet production which comprises a compound of Formula (I) and a pharmaceutically acceptable carrier.

No unacceptable toxicological effects are expected when compounds of the invention are administered in accordance with the present invention.

10 In addition, the pharmaceutically active compounds of the present invention can be co-administered with further active ingredients, such as other compounds known to treat thrombocytopenia, including chemotherapy-induced thrombocytopenia and bone marrow transplantation and other conditions with depressed platelet production, or compounds known to have utility when used in combination with a TPO mimetic.

15 Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative and not a limitation of the scope of the present invention in any way.

#### Experimental Details

20

##### Example 1

##### 3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one

25 A solution of 3,5-dibromo-2-hydroxybenzaldehyde (104 mg, 0.371 mmol) in methanol (1 mL) was added to a solution of 3-aminorhodanine (50 mg, 0.337 mmol) in methanol (5 mL) and the mixture allowed to stand at room temperature. After 1 h, the precipitate was filtered, washed (methanol, ether) and dried to give the title compound (93 mg, 67%) as a pale yellow solid. LCMS m/e 409, 411, 413 [M+H]<sup>+</sup>.

##### Example 2

30 3-(3-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1H-pyrazol-4-yl)methylene]amino]-2-thioxoimidazolidin-4-one

a) 1-(3,4-Dimethylphenyl)-3-methyl-3-pyrazolin-5-one.

A solution of 3,4-dimethylphenylhydrazine (7.3 g; 0.053 mol.) and ethyl acetoacetate (6.9 g; 0.053 mol.) in glacial acetic acid (50.0 mL) was stirred and heated at 100° for 24h. The solvent was evaporated and the product purified by chromatography (silica gel, 50% ethyl acetate/hexanes) to afford the title compound (16.8 g; 64%). MS(ES) m/z 203 [M+H].

5

b) 1-(3,4-Dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazole-4-carbaldehyde.

Phosphorus oxychloride (4.82 mL, 51.6 mmol) was added dropwise to an ice-cooled, stirred suspension of 1-(3,4-dimethylphenyl)-3-methyl-3-pyrazolin-5-one (8.70 g, 43.0 mmol) in dimethylformamide (18.0 mL) at such a rate as to maintain the temperature below 20 °C. After the addition, the mixture was heated at 100 °C for 2h, then cooled, poured into iced water (200 mL). The resulting mixture was stirred for 18h, then filtered. The solid was washed with water and dried to give the title compound (7.83 g, 79%) as a cream-coloured powder. MS (ES) m/e 231 [M+H]<sup>+</sup>.

10

15 c) 3-(3-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one.

Ethyl hydrazinoacetate hydrochloride (155 mg, 1.00 mmol) was added to a stirred solution of 3-isothiocyanatobenzoic acid (179 mg, 1.00 mmol) and di-isopropylethylamine (523  $\mu$ L, 3.00 mmol) in dichloromethane (4 mL). The mixture was stirred for 96h, evaporated under reduced pressure and partitioned between aqueous acetic acid and ethyl acetate. The organic extracts were washed with water, saturated aqueous sodium chloride, dried (magnesium sulfate) and evaporated under reduced pressure. The residue was chromatographed (silica gel, 5-15% methanol/ethyl acetate, then 20% methanol/ethyl acetate + 0.5% acetic acid) to give 1-amino-3-(3-carboxyphenyl)-2-thioxoimidazolidin-4-one (50 mg, 51%) contaminated with 6% uncyclised by-product, suitable for the next step. A solution of 1-amino-3-(3-carboxyphenyl)-2-thioxoimidazolidin-4-one (50 mg, 0.199 mmol) and 1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazole-4-carbaldehyde (55 mg, 0.239 mmol) in ethanol/methanol (2:1, 15 mL) was allowed to stand a room temperature for 96 h. The solid was filtered off, washed with ether and dried to give the title compound (42 mg, 46%) as a powder. LCMS, m/e 464 [M+H]<sup>+</sup>.

20

25

30

### Example 3

35

3-(4-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one

The procedure of example 2(c) was followed here using 4-isothiocyanatobenzoic acid instead of 3-isothiocyanatobenzoic acid to give the title compound as a powder. LCMS, m/e 464 [M+H]<sup>+</sup>.

5

#### Example 4

#### 5-(4-carboxybenzylidene)-3-[(1-{3,4-dimethylphenyl}-4-hydroxy-3-methyl-1H-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one

A mixture of 3-aminorhodanine (148 mg, 1.00 mmol), 1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1H-pyrazole-4-carbaldehyde (230 mg, 1.00 mmol) and ethanol (10 mL) was stirred 96 h. The solid was filtered, washed with ethanol and ether and dried. A mixture of the resulting crude imine (80 mg, 0.222 mmol), piperidine (2 mg, 0.022 mmol), 4-formylbenzoic acid (33 mg, 0.222 mmol), benzoic acid (3 mg, 0.022 mmol) and toluene (10 mL) was heated under reflux for 6 h in an apparatus fitted with a Dean and Stark separator to remove water. After cooling, the solid was filtered off, washed with toluene and ether, and purified by reverse phase HPLC (CombiPrep ODS-A, 10-90% acetonitrile/water + 0.1% trifluoroacetic acid) to give the title compound (18 mg, 16%) as a solid. LCMS, m/e 493 [M+H]<sup>+</sup>.

10

15

#### Example 5

#### 20 5-(3-carboxybenzylidene)-3-[(1-{3,4-dimethylphenyl}-4-hydroxy-3-methyl-1H-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one

The procedure described in example 4 was followed here using 3-formylbenzoic acid instead of 4-formylbenzoic acid to give the title compound as a powder. LCMS, m/e 493 [M+H]<sup>+</sup>.

25

#### Example 6 - Capsule Composition

An oral dosage form for administering a presently invented agonist of the TPO receptor is produced by filling a standard two piece hard gelatin capsule with the ingredients in the proportions shown in Table I, below.

30

#### Table I

<u>INGREDIENTS</u>	<u>AMOUNTS</u>
3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one (Compound A)	25 mg
Lactose	55 mg
Talc	16 mg
Magnesium Stearate	4 mg

#### Example 7 - Injectable Parenteral Composition

5 An injectable form for administering a presently invented agonist of the TPO receptor is produced by stirring 1.5% by weight of 3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one (Compound A), monosodium salt (Compound 2) in 10% by volume propylene glycol in water.

#### Example 8 - Tablet Composition

10

The sucrose, calcium sulfate dihydrate and a presently invented agonist of the TPO receptor, as shown in Table II below, are mixed and granulated in the proportions shown with a 10% gelatin solution. The wet granules are screened, dried, mixed with the starch, talc and stearic acid, screened and compressed into a tablet.

15

Table II


<u>INGREDIENTS</u>	<u>AMOUNTS</u>
3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one (Compound A)	20 mg
calcium sulfate dihydrate	30 mg
sucrose	4 mg
starch	2 mg
talc	1 mg
stearic acid	0.5 mg

20 While the preferred embodiments of the invention are illustrated by the above, it is to be understood that the invention is not limited to the precise instructions herein disclosed and that the right to all modifications coming within the scope of the following claims is reserved.



What is claimed is:



1. A method of treating of thrombocytopenia in a mammal, including a human, in need thereof which comprises administering to such mammal a therapeutically effective amount of a compound of Formula (I):



wherein:

- R<sup>1</sup> and R<sup>2</sup> are each independently selected from hydrogen, C<sub>1</sub>-12alkyl, aryl, substituted aryl, and alkyl substituted with one or more substituents selected from the group consisting of: alkoxy, acyloxy, aryl, substituted aryl, amino, N-acylamino, oxo, hydroxy, cycloalkyl, substituted cycloalkyl, -C(O)OR<sup>7</sup>, -S(O)<sub>2</sub>NR<sup>7</sup>R<sup>8</sup>, -S(O)<sub>n</sub>R<sup>6</sup>, aryloxy, nitro, cyano, halogen, and protected -OH, where
- R<sup>6</sup> is selected from hydrogen, alkyl, cycloalkyl, C<sub>1</sub>-C<sub>12</sub>aryl, substituted alkyl, substituted cycloalkyl and substituted C<sub>1</sub>-C<sub>12</sub>aryl, and
- R<sup>7</sup> and R<sup>8</sup> are independently selected from hydrogen, cycloalkyl, aryl, substituted cycloalkyl, substituted aryl, alkyl or alkyl substituted with one or more substituents selected from the group consisting of: alkoxy, acyloxy, aryloxy, amino, N-acylamino, oxo, hydroxy, -C(O)OR<sup>6</sup>, -S(O)<sub>n</sub>R<sup>6</sup>, -C(O)NR<sup>6</sup>R<sup>6</sup>, -S(O)<sub>2</sub>NR<sup>6</sup>R<sup>6</sup>, nitro, cyano, cycloalkyl, substituted cycloalkyl, halogen, C<sub>1</sub>-C<sub>12</sub>aryl, substituted C<sub>1</sub>-C<sub>12</sub>aryl and protected -OH where R<sup>6</sup> is as described above; and

- n is 0-3; or R<sup>1</sup> and R<sup>2</sup> taken together with the attached represent a ring of formula (A):




(A)

where  $R^4$  and  $R^{10}$  are each independently selected from two hydrogens,  $=NR^5$ ,  $=O$ ,  $=S$ , and  $=CHR^5$ , where  $R^5$  is  $C_1$ - $C_{12}$ aryl or substituted  $C_1$ - $C_{12}$ aryl; and  $m$  is 0 to 2;

$Z$  is a bond or selected from  $S$  or  $NR^5$ , where  $R^5$  is  $C_1$ - $C_{12}$ aryl or substituted  $C_1$ - $C_{12}$ aryl;

$R^3$  is selected from hydrogen,  $C_1$ - $C_{10}$ alkyl, phenyl, substituted phenyl, carboxyl or  $C_1$ - $C_{10}$ alkoxycarbonyl;

$L$  is a group of formula (L):



(L)

where  $A$ ,  $B$ ,  $D$  and  $E$  independently represent  $CR^{11}$  or  $N$ ; where  $R^{11}$  is selected from hydrogen, halogen,  $-CF_3$ ,  $-CN$ ,  $-SO_3H$ ,  $-SO_3Na$ ,  $-SO_2R^{14}$ ,  $-NO_2$ , phenyl, substituted phenyl,  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_{10}$ alkoxy,  $C_1$ - $C_{10}$ acyloxy, arylalkoxy,  $-COR^{14}$ ,  $-NR^{12}R^{13}$ , hydroxy or cycloalkyl;

where  $R^{14}$  is selected from hydroxy,  $C_1$ - $C_{10}$ alkyl, phenyl, amino, mono- or dialkylamino;

$R^{12}$  and  $R^{13}$  are independently selected from hydrogen,  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_{12}$ aryl, substituted  $C_1$ - $C_{12}$ aryl,  $C_1$ - $C_{10}$ acyl or cycloalkyl;

or either  $A=B$  or  $D=E$  alternatively represent  $O$ ,  $S$  or  $NR^{12}$ ; where  $R^{12}$  is as defined above;

$Y$  is selected from  $S$ ,  $O$  and  $NR^{15}$ , where  $R^{15}$  is selected from hydrogen,  $C_1$ - $C_{10}$ alkyl, substituted  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_6$ alkylphenyl, substituted  $C_1$ - $C_6$ alkylphenyl,  $C_1$ - $C_{10}$ acyl, substituted  $C_1$ - $C_{10}$ acyl, or  $SO_2R^9$ , where  $R^9$  is  $C_1$ - $C_{10}$ alkyl, substituted  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_{12}$ aryl or substituted  $C_1$ - $C_{12}$ aryl; and

$X$  is selected from  $SR^{16}$ ,  $OR^{16}$  or  $NHR^{17}$ ; where  $R^{16}$  is hydrogen,  $C_1$ - $C_{10}$ alkyl or substituted  $C_1$ - $C_{10}$ alkyl;

$R^{17}$  is hydrogen,  $C_1$ - $C_{10}$ alkyl, substituted  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_6$ alkylphenyl,  $C_1$ - $C_{10}$ acyl, substituted  $C_1$ - $C_{10}$ acyl or  $SO_2R^9$ ; where  $R^9$  is  $C_1$ - $C_{10}$ alkyl, substituted  $C_1$ - $C_{10}$ alkyl,  $C_1$ - $C_{12}$ aryl or substituted  $C_1$ - $C_{12}$ aryl; and

5 pharmaceutically acceptable salts, hydrates, solvates and esters thereof,

provided that;

10 when  $R^1$  and  $R^2$  do not form a ring and  $X$  is not  $-NHSO_2R^9$ ,  $R^5$  is not a substituted or unsubstituted pyridyl or a substituted or unsubstituted phenyl.

2. The method of claim 1 wherein the compound is selected form:

3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one;  
3-(3-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one;  
15 3-(4-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one;  
5-(4-carboxybenzylidene)-3-[(1-(3,4-dimethylphenyl)-4-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one; and  
20 5-(3-carboxybenzylidene)-3-[(1-(3,4-dimethylphenyl)-4-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one;  
or pharmaceutically acceptable salts, hydrates, solvates and esters thereof.

25 3. A method of enhancing platelet production in a mammal, including a human, in need thereof which comprises administering to such mammal a therapeutically effective amount of a compound of Claim 1.

4. The method of claim 3 wherein the compound is selected from:

30 3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one;  
3-(3-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one;  
3-(4-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one;  
5-(4-carboxybenzylidene)-3-[(1-(3,4-dimethylphenyl)-4-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one; and  
35 5-(3-carboxybenzylidene)-3-[(1-(3,4-dimethylphenyl)-4-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one;

or pharmaceutically acceptable salts, hydrates, solvates and esters thereof.

5. A pharmaceutical composition for use in enhancing platelet production which comprises a compound of Claim 1 and a pharmaceutically acceptable carrier.

5

6. Use of a compound of Formula (I), as described in claim 1, in the manufacture of a medicament for use in treating of thrombocytopenia.

7. The method of claim 1 wherein the compound is administered orally.

10

8. The method of claim 1 wherein the compound is administered parenterally.

9. A method of agonizing the TPO receptor in a subject which comprises administering an effective amount of a compound of Formula (I), as described in claim 1.

15

10. A compound represented by Formula (I) as described in claim 1.

20

11. The method of claim 1 wherein the compound is 3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one; or pharmaceutically acceptable salts, hydrates, solvates and esters thereof.

12. A compound of claim 10 selected from:

25

3-[(2-hydroxy-3,5-dibromophen-1-yl)methyleneamino]-2-thioxothiazolidin-4-one; 3-(3-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one;

3-(4-carboxyphenyl)-1-[(1-(3,4-dimethylphenyl)-5-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxoimidazolidin-4-one;

30

5-(4-carboxybenzylidene)-3-[(1-{3,4-dimethylphenyl}-4-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one; and

5-(3-carboxybenzylidene)-3-[(1-{3,4-dimethylphenyl}-4-hydroxy-3-methyl-1*H*-pyrazol-4-ylmethylene)amino]-2-thioxothiazolidin-4-one;

or a pharmaceutically acceptable salt, hydrate, solvate or ester thereof.

# INTERNATIONAL SEARCH REPORT

Inte Jonal Application No

PCT/US 00/30383

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D277/36 C07D417/12 C07D403/12 A61K31/426 A61K31/427  
A61K31/4178 A61P7/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 11262 A (ROCHE DIAGNOSTICS GMBH) 11 March 1999 (1999-03-11) the whole document	1-12
X	BECKERT R ET AL: "Zur Reaktion von Derivaten des Thiosemicarbazids mit Bisimidchloriden der Oxalsäure" MONATSHFTE FÜR CHEMIE, vol. 120, no. 12, December 1989 (1989-12), pages 1125-1137, XP002159340 the whole document, particularly page 1131, compound 11d --- -/-	10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

5 February 2001

Date of mailing of the international search report

23/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Allard, M

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 00/30383

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CROSSFIRE 'Online!  Beilstein Institut zur Foerderung der  Chemischen Wissenschaften;  XP002159341  Beilstein Registry Number 8024644, 8029346  &amp; BOLL. CHIM. FARM.,  vol. 137, no. 6, 1998, pages 210-217,</p>	5,10
X	<p>DATABASE CROSSFIRE 'Online!  Beilstein Institut zur Foerderung der  Chemischen Wissenschaften;  XP002159342  Beilstein Registry Number 217532, 1017055,  1598049, 4942046  &amp; KHIM. GETEROTSIKL. SOEDIN,  vol. 7, 1971, pages 1182-1185,</p>	5,10
X	<p>DATABASE CROSSFIRE 'Online!  Beilstein Institut zur Foerderung der  Chemischen Wissenschaften;  XP002159343  Beilstein Registry Number 1005077  &amp; LIET. TSR MOKSLU AKAD. DARB. SER. B,  1973, page 95,98</p>	10
X	<p>DATABASE CROSSFIRE 'Online!  Beilstein Institut zur Foerderung der  Chemischen Wissenschaften;  XP002159344  Beilstein Registry Number 865368, 1029545  &amp; FARM. ZH. (KIEV),  vol. 23, no. 5, 1968, pages 40-44,</p>	10
X	<p>DATABASE CROSSFIRE 'Online!  Beilstein Institut zur Foerderung der  Chemischen Wissenschaften;  XP002159345  Beilstein Registry Nimber 1088006  &amp; SB. NAUCHN. RAB., L'VOV. GOS. MED.  INST.,  vol. 24, 1963, page 22</p>	10

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

Claims Nos.: 5, 10

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty of the above mentioned claims. So many documents were retrieved that it is impossible to determine which parts of those claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT).

For these reasons it appears impossible to execute a meaningful search and/or to issue a complete search report over the whole breadth of the above mentioned claims.

The search and the report for those claims can only be considered complete for the compounds recited in claim 12.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/30383

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9911262 A	11-03-1999	AU 9265698 A	22-03-1999